

## **Appendix C — Standard Operating Procedures for Population-level Fertility Control Treatments**

### **22-month time-release pelleted vaccine:**

1. PZP vaccine would be administered only by trained BLM personnel or collaborating research partners.
2. Mares that have never been treated would receive 0.5 cc of PZP vaccine emulsified with 0.5 cc of Freund's Modified Adjuvant (FMA). Mares identified for re-treatment receive 0.5 cc of the PZP vaccine emulsified with 0.5 cc of Freund's Incomplete Adjuvant (FIA).
3. The fertility control drug is administered with two separate injections: (1) a liquid dose of PZP is administered using an 18-gauge needle primarily by hand injection; (2) the pellets are preloaded into a 14-gauge needle. These are delivered using a modified syringe and jabstick to inject the pellets into the gluteal muscles of the mares being returned to the range. The pellets are designed to release PZP over time similar to a time-release cold capsule.
4. Delivery of the vaccine would be by intramuscular injection into the gluteal muscles while the mare is restrained in a working chute. The primer would consist of 0.5 cc of liquid PZP emulsified with 0.5 cc of Freund's Modified Adjuvant (FMA). The pellets would be loaded into the jabstick for the second injection. With each injection, the liquid or pellets would be injected into the left hind quarters of the mare, above the imaginary line that connects the point of the hip (hook bone) and the point of the buttocks (pin bone).
5. In the future, the vaccine may be administered remotely using an approved long range darting protocol and delivery system if or when that technology is developed.
6. All treated mares would be freeze-marked on the hip or neck HMA managers to positively identify the animals as treated during routine field observation and at the time of possible removal during subsequent gathers.

### **Monitoring and Tracking of Treatments:**

1. At a minimum, estimation of population growth rates using helicopter or fixed-wing surveys would be conducted before any subsequent gather. During these surveys it is not necessary to identify which foals were born to which mares; only an estimate of population growth is needed (i.e. # of foals to # of adults).
2. Population growth rates of herds selected for intensive monitoring would be estimated every year post-treatment using helicopter or fixed-wing surveys. During these surveys it is not necessary to identify which foals were born to which mares, only an estimate of population growth is needed (i.e. # of foals to # of adults). If, during routine HMA field monitoring (on-the-ground), data describing mare to foal ratios can be collected, these data should also be shared with the NPO for possible analysis by the USGS.
3. A PZP Application Data sheet would be used by field applicators to record all pertinent data relating to identification of the mare (including photographs if mares are not freeze-marked) and date of treatment. Each applicator would submit a PZP Application Report and accompanying narrative and data sheets would be forwarded to the NPO (Reno, Nevada). A copy of the form and data sheets and any photos taken would be maintained at the field office.

4. A tracking system would be maintained by NPO detailing the quantity of PZP issued, the quantity used, disposition of any unused PZP, the number of treated mares by HMA, field office, and State along with the freeze-mark(s) applied by HMA and date.